

## MID-INFRARED SPECTRA OF POLYCYCLIC AROMATIC HYDROCARBON EMISSION IN HERBIG Ae/Be STARS

G. C. SLOAN,<sup>1</sup> L. D. KELLER,<sup>2</sup> W. J. FORREST,<sup>3</sup> E. LEIBENSPERGER,<sup>2</sup> B. SARGENT,<sup>3</sup> A. LI,<sup>4</sup> J. NAJITA,<sup>5</sup>  
D. M. WATSON,<sup>3</sup> B. R. BRANDL,<sup>6</sup> C. H. CHEN,<sup>5</sup> J. D. GREEN,<sup>3</sup> F. MARKWICK-KEMPER,<sup>7</sup>  
T. L. HERTER,<sup>1</sup> P. D'ALESSIO,<sup>8</sup> P. W. MORRIS,<sup>9</sup> D. J. BARRY,<sup>1</sup>  
P. HALL,<sup>1</sup> P. C. MYERS,<sup>10</sup> AND J. R. HOUCK<sup>1</sup>

Received 2005 February 7; accepted 2005 June 23

### ABSTRACT

We present spectra of four Herbig Ae/Be stars obtained with the Infrared Spectrograph (IRS) on the *Spitzer Space Telescope*. All four of the sources show strong emission from polycyclic aromatic hydrocarbons (PAHs), with the 6.2  $\mu\text{m}$  emission feature shifted to 6.3  $\mu\text{m}$  and the strongest C–C skeletal-mode feature occurring at 7.9  $\mu\text{m}$  instead of at 7.7  $\mu\text{m}$ , as is often seen. Remarkably, none of the four stars has silicate emission. The strength of the 7.9  $\mu\text{m}$  feature varies with respect to the 11.3  $\mu\text{m}$  feature among the sources, indicating that we have observed PAHs with a range of ionization fractions. The ionization fraction is higher for systems with hotter and brighter central stars. Two sources, HD 34282 and HD 169142, show emission features from aliphatic hydrocarbons at 6.85 and 7.25  $\mu\text{m}$ . The spectrum of HD 141569 shows a previously undetected emission feature at 12.4  $\mu\text{m}$  that may be related to the 12.7  $\mu\text{m}$  PAH feature. The spectrum of HD 135344, the coolest star in our sample, shows an unusual profile in the 7–9  $\mu\text{m}$  region, with the peak emission to the red of 8.0  $\mu\text{m}$  and no 8.6  $\mu\text{m}$  PAH feature.

*Subject headings:* infrared: stars — stars: chemically peculiar

### 1. INTRODUCTION

Herbig Ae/Be stars (HAeBe) are intermediate-mass (2–8  $M_{\odot}$ ) analogs to the roughly solar mass T Tauri stars (Herbig 1960; Strom et al. 1972; Waters & Waelkens 1998). The number of HAeBe stars in the Galaxy depends on the selection criteria, and as a result different studies have listed as few as  $\sim 50$  (Herbig's original list included 26 stars, but see Malfait et al. 1998) or as many as several hundred (The et al. 1994). One of the more conservative definitions assigns the HAeBe name to stars of spectral type A or B (or early F) that have broadened atomic emission lines in their optical spectra (hence the “e” in “Ae/Be”) and have infrared emission in excess of a purely photospheric spectral energy distribution. Using these criteria, Malfait et al. (1998) identified 45 HAeBe stars and published spectral energy distributions (SEDs) from UV to millimeter wavelengths, proposing an evolutionary scenario based on mid-infrared colors and SED structure.

Meeus et al. (2001) observed the 14 spatially unresolved sources (at all wavelengths) from the list by Malfait et al. (1998)

with the Short-Wavelength Spectrometer (SWS) on board the *Infrared Space Observatory* (ISO). They report the detection of PAHs in 7 of the 14 sources, proposing that the emitting PAHs are located in flared disks where the shadow of a thicker inner disk cannot attenuate the UV radiation. Acke & van den Ancker (2004) recently presented spectra from the SWS for the entire list of sources of Malfait et al. (1998), and they report the detection of PAH emission in many of them while mentioning the possibility that differences in the ionization fraction of the PAHs could help explain the structure in the 2–20  $\mu\text{m}$  spectra. They also noted that emission from crystalline silicates may confuse the issue. It is also clear from the SWS spectrum of 51 Oph (HD 158643; van den Ancker et al. 2001) that emission from gas near the stars may play a significant role in shaping the mid-infrared spectra of these stars and complicating detailed studies of PAH emission in that spectral range.

Habart et al. (2004) have analyzed ground-based and ISO observations of some 30 HAeBe stars and applied a model in which the material in the disk is in vertical hydrostatic equilibrium, is heated by the central star, and contains large dust grains in thermal equilibrium with the radiation field as well as small grains and PAHs that are transiently heated. They suggest that the PAH emission is primarily from the outer disks ( $R \sim 100$  AU) and that strong PAH emission is predominantly from sources that have flared disks (as previously proposed by Meeus et al. 2001).

Li & Lunine (2003) modeled ground-based spectra of HD 141569 in the 8–13 and 17–25  $\mu\text{m}$  regions using a porous, cometary-type dust model consisting of coagulated but otherwise unaltered interstellar grains along with fully ionized PAHs. They predict the shape of the 5–25  $\mu\text{m}$  spectrum, noting that there should be little emission from crystalline silicates (no more than 10% by mass fraction). Since PAHs are a significant component of the interstellar medium (ISM), important to the energy balance (e.g., Allamandola et al. 1989), and are apparently necessary to accurately model the emission in HAeBe stars, it is time

<sup>1</sup> Astronomy Department, Cornell University, 610 Space Sciences Building, Ithaca, NY 14853-6801; sloan@isc.astro.cornell.edu.

<sup>2</sup> Department of Physics, Ithaca College, Ithaca, NY 14850.

<sup>3</sup> Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627-0171.

<sup>4</sup> Department of Physics and Astronomy, University of Missouri at Columbia, Columbia, MO 65211.

<sup>5</sup> National Optical Astronomy Observatory, 950 North Cherry Avenue, Tucson, AZ 85719.

<sup>6</sup> Sterrewacht Leiden, P.O. Box 9513, 2300 RA Leiden, Netherlands.

<sup>7</sup> Astronomy Department, University of Virginia, P.O. Box 3818, Charlottesville, VA 22903.

<sup>8</sup> Centro de Radioastronomía y Astrofísica, UNAM, Apartado Postal 3-72 (Xangari), 58089 Morelia, Michoacan, Mexico.

<sup>9</sup> NASA *Herschel* Science Center, Infrared Processing and Analysis Center, California Institute of Technology, MS 100-22, Pasadena, CA 91125.

<sup>10</sup> Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138.

TABLE 1  
OBSERVING INFORMATION

Target	Spectral Type	$U(r)^a$ ( $\times 10^5$ )	IRS Campaign	Observation Date (UT)	AOR Key
HD 34282 .....	A0e	2.3	13	2004 Oct 10	3577856
HD 141569 .....	A0 Ve <sup>b</sup>	2.3	4	2004 Mar 3	3560960
HD 169142 .....	A5 Ve <sup>b</sup>	0.71	5	2004 Mar 26	3587584
HD 135344 .....	F4 Ve <sup>b</sup>	0.16	11	2004 Aug 8	3580672

<sup>a</sup>  $U(r)$  is the radiation field due to the star, in units of the interstellar radiation field in the solar neighborhood, calculated at  $r = 100$  AU.

<sup>b</sup> Revised spectral type from Dunkin et al. (1997).

to closely investigate not just their presence but the specific spectral properties of the emission features from PAHs around HAeBe stars.

High-sensitivity spectra of PAHs allowing precise band-strength measurements may provide information about the structure of disks and their physical and chemical conditions. The presence or absence of PAH features may provide clues to the physical properties of the disks in HAeBe systems, such as their degree of flaring (cf. Meeus et al. 2001), and may also serve as an evolutionary clock that tracks grain processing, since there are multiple destruction paths for PAHs in disks (intense irradiation, compaction, and incorporation in larger grains) but no clear formation path. The PAHs in disks around HAeBe stars most likely originate in the ISM from which the stars formed. As a significant interstellar component, PAHs may have condensed onto the ice mantles of dust grains in dense clouds that were then incorporated into protostellar nebulae. The PAHs emerge as free-flying molecules when the ice mantles of grains sublimate due to stellar irradiation or grain-grain collisions (Li & Lunine 2003). The origin of interstellar PAHs, however, is not well constrained. Suggested sources for interstellar PAHs include (1) formation around and ejection from carbon stars (Latter 1991), (2) shattering of carbonaceous interstellar dust, (3) photoprocessing of mantles on interstellar dust grains (Greenberg et al. 2000) by grain-grain collisions in interstellar shocks (Jones et al. 1996), or (4) in situ formation through ion-molecule reactions (Herbst 1991). It is essential to quantify the PAH features as a diagnostic in order to distinguish between these possibilities.

We have begun a spectroscopic survey in the mid-infrared (5.2–36  $\mu\text{m}$ ) of 16 HAeBe stars from the sample defined by Malfait et al. (1998), using IRS<sup>11</sup> (Houck et al. 2004) on the *Spitzer Space Telescope* (Werner et al. 2004). One of the goals of this study is to closely examine the PAH emission in the 5.2–14  $\mu\text{m}$  spectral region. We present the first results of our study with spectra and analysis for four stars that show PAH features, no silicate features, and few other solid-state features in their spectra. This has allowed a careful examination of the relative positions, strengths, and shapes of the features at 6.2, 7.7–7.9, 11.3, and 12.7  $\mu\text{m}$  without concern for the effects of silicate dust grains, either amorphous or crystalline. While *ISO* SWS and PHOT-S spectra of the four sources in our sample show PAH emission features (Meeus et al. 2001; Acke & van den Ancker 2004), the *Spitzer* IRS spectra have higher sensitivity, allowing the detailed analysis that we present here.

All four of our sources show evidence of circumstellar disks in their SEDs (Malfait et al. 1998). HD 34282 evidently has a large

Keplerian disk inferred from millimeter observations (Merin et al. 2004; Pietu et al. 2003), and the disk around HD 141569 has been imaged directly in the optical and near-infrared (Boccatelli et al. 2003; Mouillet et al. 2001; Weinberger et al. 1999; Augereau et al. 1999). HD 135344 has yet to be spatially resolved (Meeus et al. 2001), but Habart et al. (2005) have spatially resolved the PAH emission at 3.3  $\mu\text{m}$  in HD 169142.

## 2. OBSERVATIONS AND ANALYSIS

The four HAeBe targets were observed with the IRS on *Spitzer* in 2004 from March to October, as described in Table 1. This paper focuses on the observations with the Short-Low module (SL), which covers the 5.2–14  $\mu\text{m}$  region with a spectral resolving power of  $R \sim 90$ . The SL slit is 3"6 (2 pixels) across and 57" long.

We started with the flat-fielded images generated by the standard IRS data reduction pipeline (S11.0) at the *Spitzer* Science Center (SSC). We removed the background emission by differencing images with the source in the two SL apertures. When extracting spectra from the images, we used an extraction window that scales with wavelength; this method is similar to that used by the SSC pipeline. We obtained a spectrophotometric calibration by using identically extracted spectra of the standard star HR 6348 (K0 III) and spectral templates of this star prepared using the methods described by Cohen et al. (2003) and modified as described by G. C. Sloan et al. (2005, in preparation). The spectra in both nods and in both SL apertures were calibrated separately and then combined into one spectrum. This method generally results in very high precision within a single spectrum, indicated by the error bars in our plots, and an absolute photometric accuracy of  $\sim 10\%$  or better.

We have modified the S11.0 version of the wavelength calibration, as described by Sloan et al. (2005). This modification shifts the wavelength calibration in SL order 1 by 0.04  $\mu\text{m}$  to the blue. We estimate that our wavelengths are accurate to better than 0.02  $\mu\text{m}$ .

Figure 1 presents the spectra of our four HAeBe stars (with the spectrum of HD 135344 divided by 10 to fit on the same plot with the other three). The figure includes error bars, which are generally smaller than the vertical widths of the lines used to plot the spectra. These errors are the formal uncertainty in the mean of the spectra from the two nod positions in the SL slit (half the absolute value of the difference). With the notable exception of HD 141569, which appears to have a single cool component peaking at 40  $\mu\text{m}$ , the SEDs in our sample consist of a combination of a warm component peaking in the near-infrared ( $\sim 2$   $\mu\text{m}$ ) and a cooler component peaking at longer wavelengths (Malfait et al. 1998). This double-peaked structure may indicate that the disks have formed significant gaps between a warm inner disk

<sup>11</sup> The IRS was a collaborative venture between Cornell University and Ball Aerospace Corporation, funded by NASA through the Jet Propulsion Laboratory and the Ames Research Center.




FIG. 1.—Spectra of the sample of Herbig Ae/Be stars showing PAH emission features and no features from silicate dust grains in their spectra. The spectrum of HD 135344 has been divided by 10. These spectra do include error bars, although they are generally smaller than the width of the lines used to plot the data.

and a cool outer disk. The wavelength of the dip in intensity between the warm and cool components shifts slightly from star to star. Thus, the spectra are flat because the wavelength region observed by SL lies more in the dip between these two components. The continuum of HD 135344 appears blue because its dip appears toward longer wavelengths than the other sources and is therefore dominated by the warm component ( $T \sim 1000$  K) at the short-wavelength end of our spectrum.

Figure 2 illustrates how we measure the strengths of the PAH features, using HD 141569 as an example. We have used line segments as a rough estimate of the continuum under each feature. We anchor the line segment to the spectrum on either side of each feature and integrate the residual spectrum between the two anchor points to determine the total flux in the feature (in  $\text{W m}^{-2}$ ). Table 2 lists the wavelength intervals used to fit each feature. This method assumes that the PAH emission is optically thin, and it avoids dependencies on more sophisticated techniques to model the continuum. HD 135344 is the exception;




FIG. 2.—Spectrum of HD 141569 showing the linear-fitting method used to measure the equivalent flux in the PAH features. Table 2 gives the wavelengths used to anchor the line segments on either side of the PAH features.

TABLE 2  
FITTING WAVELENGTHS

Feature ( $\mu\text{m}$ )	$\lambda_{\text{blue}}$ ( $\mu\text{m}$ )	$\lambda_{\text{red}}$ ( $\mu\text{m}$ )
6.2.....	5.60–5.90	6.71–6.78
7.9.....	6.98–7.07	9.20–9.40 <sup>a</sup>
11.3.....	10.83–10.90	11.70–11.86
12.7.....	12.10–12.30	12.97–13.10

<sup>a</sup> 8.85–9.05  $\mu\text{m}$  for HD 135344.

before extracting the features using line segments, we first fit a polynomial estimate of the continuum to the spectrum between the features and then subtract it.

We have avoided the problem of separating the PAH features from other dust features, such as amorphous silicate emission at 10  $\mu\text{m}$  or crystalline silicate emission at 11.2  $\mu\text{m}$ , by choosing our sample carefully. The primary uncertainty is how much emission *below* the line segments belongs to the underlying PAH emission plateau and how much belongs to the feature we are measuring. If we are truncating flux from a PAH feature, we are doing so consistently to all the stars in the sample, and our relative measurements should remain valid.

The resulting profiles for the PAH features at 6.2, 7.7–7.9, 11.3, and 12.7  $\mu\text{m}$  appear in Figures 3 and 4. Table 3 presents the total fluxes and central wavelengths for each PAH feature in each of our sources. The nominal 7.7–7.9  $\mu\text{m}$  feature is shifted to 7.9  $\mu\text{m}$  (or farther) in our sample, so for the remainder of the paper, we refer to this feature as the 7.9  $\mu\text{m}$  feature. The reader should note that the flux labeled  $F_{7.9}$  in Table 3 and plotted in subsequent figures includes the emission of the 8.6  $\mu\text{m}$  feature as well. It could be more properly described as the total PAH flux from 7 to 9  $\mu\text{m}$ . We estimated the central wavelength of each feature by finding the wavelength at which half the emission lies to the blue and the red and by fitting Gaussians. The two methods produce similar results, but we adopt the Gaussian fits because they are less susceptible to noise and blended features.

Since ultraviolet and visible photons from the stellar photosphere excite the PAH emission (Li & Draine 2002), we have estimated integrated fluxes for each of our program stars relative to the interstellar radiation field (ISRF) in the solar neighborhood (Mathis et al. 1983). These estimates are intended only as a guide in our discussions of PAH excitation. Li & Lunine (2003) calculated the integrated flux ( $\lambda = 0.09\text{--}1.00$   $\mu\text{m}$ ) for HD 141569 as a function of distance,  $r$ , from the central star using spectral type B9.5 V and a Kurucz photospheric model. Table 1 presents the integrated fluxes of our stars for  $r = 100$  AU, calculated relative to their value for HD 141569. Following the nomenclature of Li & Lunine (2003),  $U(r) = 1$  corresponds to the ISRF in the solar neighborhood. We note that these are *not* in units of Habings or  $G_0$ , the ratio of the UV energy density to the estimate of Habing (1968), since we include the visible spectrum in addition to UV. Because the PAHs are emitting either from the surface of the disk or from the surrounding envelope, we assume no absorption between the photosphere and the PAH molecules.

### 3. DISCUSSION

#### 3.1. General Spectral Characteristics

The 7.7  $\mu\text{m}$  PAH feature consists of two main components, one at 7.65  $\mu\text{m}$  and the other at 7.85  $\mu\text{m}$  (Cohen et al. 1989; Bregman 1989; Peeters et al. 2002). The 7.65  $\mu\text{m}$  feature tends




FIG. 3.—The 6.2  $\mu\text{m}$  PAH emission feature and the 7–9  $\mu\text{m}$  PAH emission complex (with error bars), all normalized to the peak emission in the vicinity of 8  $\mu\text{m}$ . Line segments have been fit under the two emission regions and subtracted. In all four spectra, the “6.2  $\mu\text{m}$ ” feature has shifted to 6.3  $\mu\text{m}$ , and the peak emission around 8  $\mu\text{m}$  occurs at (or beyond) 7.9  $\mu\text{m}$  and not 7.7  $\mu\text{m}$ , as is often seen in other objects. The spectrum of HD 135344 shows an emission plateau from 8.0 to 8.2  $\mu\text{m}$  and is missing the 8.6  $\mu\text{m}$  feature.

to dominate the PAH spectra of reflection nebulae and H II regions in which the PAHs seem to have been heavily processed. The 7.85  $\mu\text{m}$  feature dominates in many planetary nebulae and objects evolving away from the asymptotic giant branch (AGB), where the PAHs seem to be relatively fresh and unprocessed. Peeters et al. (2002) studied 57 spectra of PAH sources obtained with the SWS, and they found that in spectra in which the 7.85  $\mu\text{m}$  component dominated, the 6.2  $\mu\text{m}$  PAH feature was shifted to 6.3  $\mu\text{m}$ . They used these two characteristics as the basis for what they called class B PAH spectra.

Of the 57 spectra studied by Peeters et al. (2002), 42 belong to class A, which shows the PAH emission features at 6.2 and 7.7  $\mu\text{m}$ . Class B includes 12 spectra, and class C, for which the peak emission in the 7–9  $\mu\text{m}$  range is shifted to the red of 8.0  $\mu\text{m}$ ,

accounts for two sources. One source in their sample had characteristics of both class A and B.

Figure 5 compares the spectra of HD 141569 and HD 135344 in our sample to some of the sources considered by Peeters et al. (2002). The top panel illustrates PAH spectra of classes A, B, and C, as defined by Peeters et al. (2002), using as prototypes IRAS 03260+3111 (a young stellar object in the NGC 1333 molecular cloud core), HD 44719, (the Red Rectangle), and AFGL 2688 (the Cygnus Egg), respectively. The comparison spectra were obtained from the SWS atlas (Sloan et al. 2003). All the IRS spectra and the SWS spectra of IRAS 03260 and HD 44719 have had the continuum removed using line segments, as illustrated in Figure 2. The SWS spectra of AFGL 2688 and HD 100546 have steep red continua; to isolate the PAH




FIG. 4.—PAH emission features at 11.3 and 12.7  $\mu\text{m}$  (with error bars), after continuum subtraction. Separate line segments have been fit under the features and removed. All spectra are normalized at 11.3  $\mu\text{m}$ . The most ionized spectrum, HD 141569, shows an emission feature at 12.4  $\mu\text{m}$ .

TABLE 3  
FEATURE POSITIONS AND STRENGTHS

TARGET	6.2 $\mu\text{m}$ FEATURE		7.9 $\mu\text{m}$ FEATURE		11.3 $\mu\text{m}$ FEATURE		12.7 $\mu\text{m}$ FEATURE	
	$\lambda$ ( $\mu\text{m}$ )	Flux ( $10^{-15} \text{ W m}^{-2}$ )	$\lambda$ ( $\mu\text{m}$ )	Flux ( $10^{-15} \text{ W m}^{-2}$ )	$\lambda$ ( $\mu\text{m}$ )	Flux ( $10^{-15} \text{ W m}^{-2}$ )	$\lambda$ ( $\mu\text{m}$ )	Flux ( $10^{-15} \text{ W m}^{-2}$ )
HD 34282 .....	6.26	$12.63 \pm 0.17$	7.92	$47.09 \pm 0.58$	11.27	$3.11 \pm 0.14$	12.64	$0.54 \pm 0.06$
HD 141569 .....	6.26	$6.84 \pm 0.15$	7.98	$16.65 \pm 0.35$	11.28	$0.62 \pm 0.10$	12.52	$0.18 \pm 0.02$
HD 169142 .....	6.26	$32.71 \pm 0.54$	8.01	$88.06 \pm 0.93$	11.30	$11.86 \pm 0.58$	12.71	$1.95 \pm 0.17$
HD 135344 .....	6.28	$3.47 \pm 0.51$	8.02	$12.49 \pm 1.30$	11.36	$1.38 \pm 0.21$	12.75	$0.18 \pm 0.06$




FIG. 5.— Comparison of spectra from the SWS on *ISO* with two spectra from our sample. *Top*: Prototypical SWS spectra of the three classes defined by Peeters et al. (2002). IRAS 03260+3111 is a young stellar object, while HD 44179 (the Red Rectangle) and AFGL 2688 (the Cygnus Egg) are post-AGB objects. *Middle*: HD 141569 compared to two class B objects. HD 141569 resembles the HAeBe star more than HD 44179. *Bottom*: HD 135344 compared to HD 100546 and the class C spectrum from AFGL 2688, showing that the spectrum resembles a blend between the two.

spectrum we fitted a polynomial to the continuum and then subtracted it.

Figure 3 and the middle panel of Figure 5 show that our HAeBe stars have spectral features typical of class B, with the  $6.2\ \mu\text{m}$  feature shifted to at least  $6.26\ \mu\text{m}$ , and the peak emission in the  $7.6\text{--}7.9\ \mu\text{m}$  region falling near  $7.9\ \mu\text{m}$ . Both the  $6.2$  and  $7.9\ \mu\text{m}$  features arise from C—C stretching modes. The  $8.6\ \mu\text{m}$  feature, which is included in our measurements of the strength of the  $7.9\ \mu\text{m}$  feature, probably arises from a C—H in-plane bending mode.

The sample examined by Peeters et al. (2002) includes two isolated HAeBe stars, HD 100546 and HD 179218, which they included with the class B PAH spectra. The middle panel of Figure 5 shows that in the spectrum of HD 100546, the  $7.9\ \mu\text{m}$  PAH feature is actually shifted to even longer wavelengths, nearly to  $8.0\ \mu\text{m}$ . In the other HAeBe star in the SWS sample, this feature appears at  $7.8\ \mu\text{m}$ , but in all four of the IRS spectra, this feature lies closer to  $8.0\ \mu\text{m}$ . Thus, in five of the six isolated HAeBe stars studied here or by Peeters et al. (2002), the  $7.8\text{--}7.9\ \mu\text{m}$  feature indicative of class B is shifted to nearly  $8.0\ \mu\text{m}$ . The HAeBe stars may represent a subgroup within class B.

Other features visible in the spectra in Figure 1 in the  $5\text{--}8\ \mu\text{m}$  range require comment as well. The  $5.2\ \mu\text{m}$  PAH emission feature appears in three of the spectra at the short-wavelength cutoff, but it is absent in HD 34282. Two of the spectra, HD 34282 and HD 169142, show spectral structure in the vicinity of  $7\ \mu\text{m}$ , which is due to emission features at  $6.85$  and  $7.25\ \mu\text{m}$ .

Chiar et al. (2000) made the first detection of absorption features at  $6.85$  and  $7.25\ \mu\text{m}$ , which they attributed to C—H stretching modes in aliphatic hydrocarbons such as methyl or methylene groups. This discovery was made in the SWS spectrum of Sgr A\*. Bouwman et al. (2001) detected the  $6.85\ \mu\text{m}$  band in emission in the HAeBe star HD 163296. Spoon et al. (2004) recently reported the  $6.85$  and  $7.25\ \mu\text{m}$  bands in absorption in the spectrum of the ultraluminous infrared galaxy IRAS F00183—7111, and they showed that laboratory data of hydrogenated amorphous carbon (HACs; Furton et al. 1999) provide a good match. We note that the  $3.4\ \mu\text{m}$  interstellar absorption band, identified as an aliphatic C—H stretching mode (Sandford et al. 1991), may also appear in emission (Geballe & van der Veen 1990; Geballe et al. 1992). It is likely therefore that we have observed these aliphatic HAC features in emission in HD 34282 and HD 169142.

### 3.2. HD 135344

The spectrum of HD 135344 contains a much stronger contribution from a warm continuum ( $T \sim 1000\ \text{K}$ ) underlying the PAHs, making the shape and position of the weaker PAH features much more susceptible to any systematic problems with the extraction and calibration of the spectrum, which explains the larger error bars in its spectrum shown in Figure 3. Nonetheless, some differences from the other spectra in the sample are still apparent.

In the  $7\text{--}9\ \mu\text{m}$  region, HD 135344 shows a plateau extending from  $8.0$  to  $8.2\ \mu\text{m}$ , most likely a blend of peaks at these two wavelengths. The  $8.0\ \mu\text{m}$  peak is similar to that seen in four of the other HAeBe stars, but the  $8.2\ \mu\text{m}$  peak more closely resembles class C PAH spectra (Peeters et al. 2002). Both of the objects in this class, AFGL 2688 and IRAS 13416—6243, are post-AGB objects, and neither shows an  $8.6\ \mu\text{m}$  feature. HD 135344 has more emission shortward of  $8.0\ \mu\text{m}$  than these two objects, but the missing  $8.6\ \mu\text{m}$  feature is striking.

The spectrum of HD 135344 differs from the other HAeBe stars in other ways. The  $11.3\ \mu\text{m}$  feature is shifted by at least

$0.06\ \mu\text{m}$  to the red; this is too large to have arisen from the random pointing errors that occasionally place a star off the central axis of the IRS slit. HD 135344 also has a relatively weak contribution at  $6.0\ \mu\text{m}$  (compared to  $6.3\ \mu\text{m}$ ).

Beintima et al. (1996) attributed the  $6.0\ \mu\text{m}$  satellite of the  $6.2\ \mu\text{m}$  feature to a weak C—C mode, but we note that in the objects studied by Peeters et al. (2002) the strength of the  $6.0\ \mu\text{m}$  feature shows no correlation with that of the  $6.2\ \mu\text{m}$  C—C mode. A correlation between the strength of the two bands also seems absent in our sample. Peeters et al. (2002) speculated that the  $6.0\ \mu\text{m}$  feature arises from C—O stretching modes in oxygenated PAH species, but these modes generally peak at somewhat shorter wavelengths (Schutte et al. 1990). The carrier of the  $6.0\ \mu\text{m}$  feature remains uncertain.

Oudmaijer et al. (1992) have revised the spectral classification of HD 135344 from A0 V (Houk 1982) to F4 Ve, and Dunkin et al. (1997) have confirmed this new spectral type.<sup>12</sup> It is interesting that this source, which is significantly cooler than the other three sources in the sample, is the one that shows the most unusual PAH emission in the  $8\ \mu\text{m}$  region.

### 3.3. Dependence of PAH Emission on Ionization Fraction

The relative strengths of PAH features can be used to identify the degree of ionization of the PAHs. They may therefore provide clues to the incident radiation field, electron density, and gas temperature. The most obvious difference between neutral and ionized PAHs is the relative strength of the C—C modes in the  $7.7\text{--}7.9\ \mu\text{m}$  region compared to the out-of-plane C—H bending modes at  $11.3\ \mu\text{m}$  and longer wavelengths (e.g., Allamandola et al. 1999). Ionized PAHs emit more strongly in the features at  $6\text{--}9\ \mu\text{m}$  than in the  $10\text{--}13\ \mu\text{m}$  region, while in neutral PAHs, the  $10\text{--}13\ \mu\text{m}$  region dominates. The models of Draine & Li (2001) show that the ratio  $F_{7.9}/F_{11.3}$  increases with the size of the PAH molecule, since larger PAHs generally have a higher C:H (e.g., see Li & Draine 2001). However, a shift from neutrals to ions increases the ratio much more significantly, by a factor of  $\sim 10$ .

We can thus use the ratio of the strengths of the  $7.9$  and  $11.3\ \mu\text{m}$  features as an estimate for the relative degree of ionization in each spectrum. This ratio is the most obvious difference among the spectra in our sample, varying from  $7.4$  in HD 169142 to  $25.2$  in HD 141569, making these the least and most ionized PAH spectra, respectively. The ratios for the intermediate stars are  $9.1$  for HD 135344 and  $15.2$  for HD 34282.

Recent optical spectroscopy by Dunkin et al. (1997) confirms the previous revision of the spectral classification of HD 135344, as discussed above, and also revises the classification of HD 169142 from B9 V (Houk 1982) to A5 Ve. These new classifications correlate with the observed ionization fractions as measured by the  $F_{7.9}/F_{11.3}$  ratio, with the two A0 stars having the more ionized spectra and the later type stars having the less ionized spectra.

In HD 141569, according to detailed models (Li & Lunine 2003), the charging of PAHs is dominated by collisions with electrons and not by emitting photoelectrons. The electron recombination rate is 3 orders of magnitude larger than the photoelectron emission rate. For this case, then, the PAHs are

<sup>12</sup> Zuckerman et al. (1995) and Coulson & Walther (1995) find even later spectral types, F6 V and F8 V, respectively. Coulson & Walther (1995) point out that HD 135344 is a visual binary with the two components, SAO 206462 and 206463, separated by  $20''$ . SAO 206463 is the brighter source and is probably the origin of the A0 V classification, but the coordinates of HD 135344 match those of the fainter SAO 206462, which is the target under consideration here.

negatively charged. Our analysis is not sensitive to the polarity of the charge of the PAHs; it is sensitive only to the ionization fraction.

### 3.4. The 7.9 $\mu\text{m}$ Feature

There is no clear explanation for the varying strengths of the components of the 7–9  $\mu\text{m}$  PAH complex among observed sources. Bregman & Temi (2005) investigate two possible causes in their study of the spatial distribution of the components in extended reflection nebulae. The peak of the complex might shift from 7.6 to 7.9  $\mu\text{m}$  as the fraction of PAH cations increases or as the fraction of fresh and unprocessed PAHs increases. Our sample shows a range of ionization fractions, but in three of the four spectra, the shape of the 7–9  $\mu\text{m}$  complex looks similar. The ratio of the 7.9 and 11.3  $\mu\text{m}$  features varies by a factor of 3.7 in these three sources, and yet the relative contributions of the 7.6 and 7.9  $\mu\text{m}$  components seem fixed, which would suggest that the components are not sensitive to ionization, at least over the ionization range we have examined.

The size of the PAH molecules may also influence the relative strengths of the features at 7.6 and 7.9  $\mu\text{m}$ , but there is some disagreement over how the position of the strongest feature in this wavelength range depends on PAH size. Boissel et al. (2001) suggest that very small grains produce the longer-wavelength feature, but Hudgins & Allamandola (1999a) report that their laboratory measurements show that the 7.6  $\mu\text{m}$  feature shifts to longer wavelength for *larger* PAHs. We can only say that our data are least consistent with the dependence of the 7–9  $\mu\text{m}$  PAH complex on the ionization fraction.

### 3.5. Possible Correlations with PAH Size and Structure

The models of Draine & Li (2001) show a dependence of the ratio  $F_{6.2}/F_{7.9}$  on the size of the PAH molecule, but it is weak, with the flux ratio doubling only when the number of C atoms has been reduced by a factor of  $\sim 10$ . The top panel of Figure 6 shows that the ratio  $F_{6.2}/F_{7.9}$  varies by a factor of 2 in our sample, but not in any systematic way with the ionization fraction as indicated by  $F_{7.9}/F_{11.3}$ .

To better discriminate the size of the PAH molecules, Hony et al. (2001) use the ratio  $F_{12.7}/F_{11.3}$ . The 11.3  $\mu\text{m}$  feature arises from a C–H stretching mode when an aromatic ring contains only one H atom, such as one might find along the edge of a large PAH molecule. The 12.7  $\mu\text{m}$  feature arises from the C–H trio stretching mode, i.e., from aromatic rings with three H atoms, which are found on the corners and ends of PAH molecules. As one increases the size of a PAH molecule, the strength of the emission at 11.3  $\mu\text{m}$  increases with respect to the emission at 12.7  $\mu\text{m}$ .

The bottom panel of Figure 6 shows that the strength of the 12.7  $\mu\text{m}$  feature generally increases with the ionization fraction. Comparing this panel with the top panel reveals that the use of the ratios  $F_{6.2}/F_{7.9}$  and  $F_{12.7}/F_{11.3}$  as size indicators gives contradictory results.

The  $F_{12.7}/F_{11.3}$  ratio may be influenced not just by the size of the PAHs but by their geometry. If a compact PAH is rearranged into a structure with the same number of carbon atoms but with more uneven edges, the strength of the duo and trio modes increases compared to the solo mode. Thus, it is possible that the enhancement of the 12.7  $\mu\text{m}$  feature could also arise from more irregularly shaped and less compact PAHs.

Measurements at the Astrochemistry Laboratory at the NASA Ames Research Center have revealed no dependency of the  $F_{6.2}/F_{7.9}$  ratio on the size of the PAHs (D. M. Hudgins 2004,




FIG. 6.—Flux ratios from the different PAH features plotted vs. the ratio of the 7.9 and 11.3  $\mu\text{m}$  PAH features. The ratio  $F_{7.9}/F_{11.3}$  traces the ionization fraction of the PAHs. From left to right (least to most ionized), the circles are for HD 169142 (A5), HD 135344 (F4), HD 34282 (A0), and HD 141569 (A0). *Top*: The relative strengths of the 6.2 and 7.9  $\mu\text{m}$  features vary by a factor of  $\sim 2$ , but they do not vary systematically with ionization fraction. *Bottom*: The ratio  $F_{12.7}/F_{11.3}$  increases with ionization fraction, suggesting that the size of the PAHs decreases with ionization fraction.

private communication), but the Astrochemistry Laboratory has not explored a full order of magnitude in PAH size. While larger PAHs such as those modeled by Draine & Li (2001) do show a dependence, typical PAH sizes are on the order of  $\sim 100$  carbon atoms (e.g., Li & Draine 2001), and the range in PAH sizes may be insufficient to dominate other causes for variation in the strength of the 6.2 and 7.9  $\mu\text{m}$  bands. The relative strengths of the out-of-plane C–H bending modes, on the other hand, are much more sensitive to variations in PAH size, especially for the sizes of the molecules likely to be encountered in typical astrophysical environments.

The spectra of our HAeBe stars show other emission features in the 11–13  $\mu\text{m}$  region in addition to the 11.3 and 12.7  $\mu\text{m}$  features. In Figures 1 and 4, the duet out-of-plane C–H bending mode at 11.9  $\mu\text{m}$  is visible in all of the spectra except HD 135344. These figures also show that what we are measuring as the 12.7  $\mu\text{m}$  feature shows some structure. In the spectrum of HD 141569, the secondary emission peak at 12.4  $\mu\text{m}$  actually dominates the 12.7  $\mu\text{m}$  feature. The Humphreys- $\alpha$  line should be at 12.365  $\mu\text{m}$ , but there is no corresponding Pfund- $\alpha$  line at 7.46  $\mu\text{m}$ . The  $\text{H}_2$  0–0  $S(2)$  is expected at 12.28  $\mu\text{m}$ , but this is too far to the blue, and it is not accompanied by the 0–0  $S(3)$  line at 9.67  $\mu\text{m}$ . The most likely carrier of the 12.4  $\mu\text{m}$  feature is another trio out-of-plane bending mode, as this feature

can vary from 12.1 to 13.5  $\mu\text{m}$ , depending on the PAH molecule (e.g., Hudgins & Allamandola 1999b).

#### 4. CONCLUSION

We have presented mid-infrared spectroscopy from *Spitzer* of four Herbig Ae/Be stars whose spectra show strong PAH emission but no silicate dust features. The absence of silicate emission has enabled a detailed analysis independent of specific disk models of the relative strengths and positions of PAH emission features in the 5–14  $\mu\text{m}$  region.

Our sample shows spectral characteristics generally consistent with the class B PAH emission sources identified by Peeters et al. (2002), viz., a 6.2  $\mu\text{m}$  feature shifted to 6.3  $\mu\text{m}$  and 7–9  $\mu\text{m}$  emission dominated by the component at 7.9  $\mu\text{m}$ . However, in all of our sources, the 7.9  $\mu\text{m}$  feature is shifted closer to 8.0  $\mu\text{m}$ , as in the spectrum of HD 100546, a HAeBe star observed by the SWS on *ISO*.

The spectra show a range in ionization fraction as revealed by the  $F_{7.9}/F_{11.3}$  ratio, with HD 141569 showing the highest degree of ionization and HD 169142 the lowest. The ionization fraction is lower for later spectral classes.

The strength of the 12.7  $\mu\text{m}$  feature increases with respect to the 11.3  $\mu\text{m}$  feature as the ionization fraction increases. The strength of the 6.2  $\mu\text{m}$  feature relative to the 7.9  $\mu\text{m}$  feature varies by a factor of nearly 2 in our sample, but in no systematic way.

The spectrum of HD 135344, the coolest star in our sample, resembles the class C spectra discussed by Peeters et al. (2002) in some ways, with more emission to the red of 7.9  $\mu\text{m}$  than in the other sources and a missing 8.6  $\mu\text{m}$  feature.

Thus, our sample of PAH spectra from HAeBe stars shows some characteristics that reinforce results from the literature but also raise new questions. Why do HAeBe stars consistently produce the class B PAH spectra seen in less than 25% of the SWS sample? What causes the shift of the 7.9  $\mu\text{m}$  feature to  $\sim$ 8.0  $\mu\text{m}$ ? What causes the range of 6.2  $\mu\text{m}$  feature strengths compared to the 7.9  $\mu\text{m}$  feature? What is the carrier of the 6.0  $\mu\text{m}$  feature? Answering these questions will help us understand the nature of the HAeBe systems from which the PAH emission arises.

Quantitative measures of the size and ionization fraction of the PAHs would be valuable in determining the degree of grain processing in the disks and the fraction of short-wavelength stellar photons that reach the outer disks. With the small size of the current sample, it is not possible to determine the origin of the correlations, although we expect that the full IRS sample of HAeBe stars will bring us closer to this goal.

J. D. Bregman, D. M. Hudgins, and L. J. Allamandola provided useful comments as we prepared this manuscript. This work is based on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under NASA contract 1407. Support for this work was provided by NASA through contract 1257184 issued by JPL/Caltech. L. D. K. was supported in part by awards from the Research Corporation and from NASA, administered by the American Astronomical Society. This research has made use of the SIMBAD database operated at the Centre de Données astronomiques de Strasbourg.

#### REFERENCES

- Acke, B., & van den Ancker, M. E. 2004, *A&A*, 426, 151  
 Allamandola, L. J., Hudgins, D. M., & Sandford, S. A. 1999, *ApJ*, 511, L115  
 Allamandola, L. J., Tielens, A. G. G. M., & Barker, J. R. 1989, *ApJS*, 71, 733  
 Augereau, J. C., Lagrange, A. M., Mouillet, D., & Menard, F. 1999, *A&A*, 350, L51  
 Beintima, D. A., et al. 1996, *A&A*, 315, L369  
 Bocatelli, A., Augereau, J. C., Marchis, F., & Hahn, J. 2003, *ApJ*, 585, 494  
 Boissel, P., Joblin, C., & Pernot, P. 2001, *A&A*, 373, L5  
 Bouwman, J., Meeus, G., de Koter, A., Hony, S., Dominik, C., & Waters, L. B. F. M. 2001, *A&A*, 375, 950  
 Bregman, J. D. 1989, in *IAU Symp. 135, Interstellar Dust*, ed. L. J. Allamandola & A. G. G. M. Tielens (Dordrecht: Kluwer), 109  
 Bregman, J. D., & Temi, P. 2005, *ApJ*, 621, 831  
 Chiar, J. E., Tielens, A. G. G. M., Whittet, D. C. B., Schutte, W. A., Boogert, A. C. A., Lutz, D., van Dishoeck, E. F., & Bernstein, M. P. 2000, *ApJ*, 537, 749  
 Cohen, M., Megeath, T. G., Hammersley, P. L., Martin-Luis, F., & Stauffer, J. 2003, *AJ*, 125, 2645  
 Cohen, M., Tielens, A. G. G. M., Bregman, J., Witteborn, F. C., Rank, D. M., Allamandola, L. J., Wooden, D., & Jourdain de Muizon, M. 1989, *ApJ*, 341, 246  
 Coulson, I. M., & Walther, D. M. 1995, *MNRAS*, 274, 977  
 Draine, B. T., & Li, A. 2001, *ApJ*, 551, 807  
 Dunkin, S. K., Barlow, M. J., & Ryan, S. G. 1997, *MNRAS*, 286, 604  
 Furton, D. G., Laiho, J. W., & Witt, A. N. 1999, *ApJ*, 526, 752  
 Geballe, T. R., Tielens, A. G. G. M., Kwok, S., & Hrivnak, B. J. 1992, *ApJ*, 387, L89  
 Geballe, T. R., & van der Veen, W. E. C. J. 1990, *A&A*, 235, L9  
 Greenberg, J. M., et al. 2000, *ApJ*, 531, L71  
 Habart, E., Natta, A., & Krugel, E. 2004, *A&A*, 427, 179  
 Habart, E., Natta, A., Testi, L., & Carillet, M. 2005, *A&A*, in press  
 Habing, H. J. 1968, *Bull. Astron. Inst. Netherlands*, 19, 421  
 Herbig, G. H. 1960, *ApJS*, 4, 337  
 Herbst, E. 1991, *ApJ*, 366, 133  
 Hony, S., Van Kerckhoven, C., Peeters, E., Tielens, A. G. G. M., & Allamandola, L. J. 2001, *A&A*, 370, 1030  
 Houck, J. R., et al. 2004, *ApJS*, 154, 18  
 Houk, N. 1982, *Michigan Catalogue of Two-Dimensional Spectral Types for the HD Stars*, Vol. 3 (Ann Arbor: Univ. Michigan)
- Hudgins, D. M., & Allamandola, L. J. 1999a, *ApJ*, 513, L69  
 ———. 1999b, *ApJ*, 516, L41  
 Jones, A. P., Tielens, A. G. G. M., & Hollenbach, D. J. 1996, *ApJ*, 469, 740  
 Latter, W. B. 1991, *ApJ*, 377, 187  
 Li, A., & Draine, B. T. 2001, *ApJ*, 554, 778  
 ———. 2002, *ApJ*, 576, 762  
 Li, A., & Lunine, J. I. 2003, *ApJ*, 594, 987  
 Malfait, K., Bogaert, E., & Waelkens, C. 1998, *A&A*, 331, 211  
 Mathis, J. S., Mezger, P. G., & Panagia, N. 1983, *A&A*, 128, 212  
 Meeus, G., Waters, L. B. F. M., Bouwman, J., van den Ancker, M. E., Waelkens, C., & Malfait, K. 2001, *A&A*, 365, 476  
 Merin, B., et al. 2004, *A&A*, 419, 301  
 Mouillet, D., Lagrange, A. M., Augereau, J. C., & Menard, F. 2001, *A&A*, 372, L61  
 Oudmaijer, R. D., van der Veen, W. E. C. J., Waters, L. B. F. M., Trams, N. R., Waelkens, C., & Engelsman, E. 1992, *A&AS*, 96, 625  
 Peeters, E., Hony, S., Van Kerckhoven, C., Tielens, A. G. G. M., Allamandola, L. J., Hudgins, D. M., & Bauschlicher, C. W. 2002, *A&A*, 390, 1089  
 Pietu, V., Dutrey, A., & Kahane, C. 2003, *A&A*, 398, 565  
 Sandford, S. A., Allamandola, L. J., Tielens, A. G. G. M., Sellgren, K., Tapia, M., & Pendleton, Y. 1991, *ApJ*, 371, 607  
 Schutte, W. A., Tielens, A. G. G. M., Allamandola, L. J., Wooden, D. H., & Cohen, M. 1990, *ApJ*, 360, 577  
 Sloan, G. C., Kraemer, K. E., & Price, S. D. 2003, *ApJS*, 147, 379  
 Sloan, G. C., Spoon, H. W. W., & Bernard-Salas, J. 2005, *Improving the Wavelength Calibration of Short-Low and Long-Low (IRS Tech. Rep. 05002; Ithaca: Cornell)*, <http://isc.astro.cornell.edu/tech/tr>  
 Spoon, H. W. W., et al. 2004, *ApJS*, 154, 184  
 Strom, S. E., Strom, K. M., Yost, J., Carrasco, L., & Grasdalen, G. 1972, *ApJ*, 173, 353  
 The, P. S., de Winter, D., & Perez, M. R. 1994, *A&AS*, 104, 315  
 van den Ancker, M. E., Meeus, G., Cami, J., Waters, L. B. F. M., & Waelkens, C. 2001, *A&A*, 369, L17  
 Waters, L. B. F. M., & Waelkens, C. 1998, *ARA&A*, 36, 233  
 Weinberger, A. J., et al. 1999, *ApJ*, 525, L53  
 Werner, M. W., et al. 2004, *ApJS*, 154, 1  
 Zuckerman, B., Forveille, T., & Kastner, J. H. 1995, *Nature*, 373, 494